

Processes for the detection of nucleic acidsPatent Number: ☐ US5344757

Publication date: 1994-09-06

Inventor(s): KESSLER CHRISTOPH (DE); MATTES RALF (DE); SCHMITZ GUDRUN (DE);
SCHOELER HANS R (DE); SEIBL RUDOLF (DE); HOELTKE HANS-JOACHIM (DE)

Applicant(s): BOEHRINGER MANNHEIM GMBH (DE)

Requested
Patent: ☐ EP0324474, B1Application
Number: US19900415307 19900109Priority Number
(s): DE19883800642 19880112; DE19883813278 19880420; WO1989EP00026 19890112IPC
Classification: C12Q1/68EC
Classification: C07H21/00G, C12Q1/68B2, C12Q1/68A2Equivalents: ☐ DE3813278, ES2054883T, HK116996, JP1503647T, JP7031194B, ☐ WO8906698

Abstract

PCT No. PCT/EP89/00026 Sec. 371 Date Jan. 9, 1990 Sec. 102(e) Date Jan. 9, 1990 PCT Filed Jan. 12, 1989 PCT Pub. No. WO89/06698 PCT Pub. Date Jul. 27, 1989. For the detection of nucleic acids of definite sequence by hybridisation with a complementary nucleic acid probe which contains bound via a chemical bonding at least one hapten as labelling one uses, as hapten, a steroid which is bound via a bridge of at least 4 atoms length to at least one position of the nucleic acid probe which does not participate in hydrogen bridge formation and detects the hybridised probe via an in turn labelled anti-hapten antibody.

Data supplied from the esp@cenet database - I2